

**The Higher Risk General Surgical Patient :
Towards Improved Care for a Forgotten Group**

Report

of the

**Royal College of Surgeons of England /
Department of Health
Working Group**

on

**Peri-operative Care
of the
Higher-Risk General Surgical Patient**

(Final Draft)

Contents

| | |
|--|----|
| Contributors | 3 |
| Approving Organisations | 3 |
| Summary | 4 |
| Key Recommendations | 5 |
| Background | 7 |
| Introduction | 7 |
| Variation in Outcomes | 9 |
| How do adverse outcomes occur for the higher-risk general surgery patient? | 10 |
| Sepsis | 12 |
| Actions | 14 |
| Managing the critically ill surgical patient with sepsis | 14 |
| Escalation of care | 14 |
| Urgency of source control | 16 |
| Summary timelines | 18 |
| Assessing and identifying risk | 19 |
| End of Surgery Bundle | 22 |
| Post-Operative Care | 24 |
| Structured care on the Post anaesthetic care unit (PACU) | 25 |
| Co-location of medium risk patients | 26 |
| Audit and outcomes | 27 |
| Conclusions | 28 |
| Appendix 1 Early Warning Score (EWS) | 30 |
| Appendix 2 Pathway for response to the unwell surgical patient | 31 |
| References | 34 |

Royal College of Surgeons of England / Department of Health Working Group on Peri-operative Care of the Higher-Risk General Surgical Patient

I D Anderson Consultant General & Colorectal Surgeon, Salford Royal Foundation NHS Trust (*Chair, RCS, ASGBI*)

J Eddleston Consultant Anaesthetist and Intensivist, Central Manchester Foundation NHS Trust (*Adult Critical Care Advisor Department of Health*)

M Grocott Consultant Anaesthetist and Intensivist, Southampton, Director, NIAA Health Services Research Centre, The Royal College of Anaesthetists (*RCoA*)

N P Lees Consultant General & Colorectal Surgeon, Salford Royal Foundation NHS Trust (*ACPGBI*)

D Lobo Consultant General & Upper GI Surgeon, Queens Medical Centre, Nottingham. (*AUGIS, SARS*)

I Loftus Consultant Vascular Surgeon, St George's Hospital, London. (*RCS, VS*)

N Markham Consultant General Surgeon, North Devon. (*ASGBI*)

D Mitchell Consultant Vascular and Renal Transplant Surgeon
Chair, Audit and Quality Improvement Committee. Vascular Society of Great Britain & Ireland (*VS*)

R Pearse Senior Lecturer & Consultant in Intensive Care Medicine, Barts and The London School of Medicine and Dentistry.

C Peden Consultant Anaesthetist and Intensivist, Royal United Hospital, Bath.

R D Sayers Professor of Vascular Surgery, University of Leicester. (*RCS, VS*)

J Wigfull Consultant Anaesthetist and Intensivist, Sheffield Teaching Hospitals.

Approving Organisations

This document has been approved by

| | |
|--------|--|
| ASGBI | Association of Surgeons of Great Britain and Ireland |
| ACPGBI | Association of Coloproctology of Great Britain and Ireland |
| AUGIS | Association of Upper Gastro-intestinal Surgeons |
| SARS | Society of Academic and Research Surgeons |

Summary

Higher-risk non-cardiac general surgery is undertaken in every acute hospital. By way of comparison, the mortality for this group, which includes most major gastro-intestinal and vascular procedures, exceeds that for cardiac surgery by two to three fold and complication rates of 50% are not uncommon. There may be a lack of awareness of the level of risk. Among these patients, emergency surgery and unscheduled management of complications is common and this group of patients are one of the largest consumers of critical care resources. The health and financial costs are considerable.

Evidence indicates that the peri-operative pathway followed by patients requiring emergency management is frequently disjointed, protracted and not always patient centred. Outcomes are known to vary substantially and could be considerably improved. Patients at higher risk can be identified and should receive differential management as recommended below.

It is the opinion of this expert group that the recommendations contained within should be deliverable within 2 years in all acute hospitals undertaking complex or unscheduled general surgery in adults and that doing so would make an appreciable difference to outcomes.

Key recommendations

1. Trusts should formalise their pathways for unscheduled adult general surgical care. All patients should have a clear diagnostic and monitoring plan on admission. The monitoring plan must be compliant with *NICE 50* guidance and match competency of the doctor and nurse to needs of the patient.
2. Clinical management must reflect the estimated risk with due adjustments being made in the seniority of staff involved, pre-operative preparation, urgency of treatment and utilisation of critical care. This is of particular importance during unscheduled care which accounts for 80% or more of postoperative deaths and complications. Differing structures may be needed for elective and emergency cases.
3. Delay in interventional treatments is currently not uncommon and leads to avoidable death and complications. The urgency of treatment should match the severity of the patient's condition as specified in this document. Hospitals should provide adequate emergency theatre access and prioritise emergencies ahead of elective work whenever necessary.
4. Each patient should have their expected risk of death identified and documented prior to surgery or other intervention. High risk patients are defined by a predicted hospital mortality of $\geq 10\%$.
5. Surgical procedures with a predicted mortality of $\geq 10\%$ should be conducted under the direct supervision of a consultant surgeon and a consultant anaesthetist.
6. Analogous considerations apply to staffing, seniority, urgency and support when radiological intervention is planned.
7. A review of the progress of higher risk operations should be conducted jointly by surgeon and anaesthetist towards the end of the procedure to determine the optimal location of further management. Pathways for post-surgical care must be developed based on clinical risk of deterioration and other factors such as those within the *End of Surgery* bundle described. Patients with an estimated risk of death $\geq 10\%$ must be admitted to a critical care location.

8. Prompt recognition and treatment of complications is essential to minimise avoidable morbidity and mortality and hence reduce costs. These requirements apply particularly to surgical patients with sepsis who often require complex management and who are amongst the largest users of critical care resources. Delay in the management of these patients worsens outcomes. The adoption of an escalation strategy, which incorporates defined time-points and the early involvement of senior staff when necessary is strongly advised. One such strategy is defined.
9. The heterogeneity and urgent nature of much higher-risk surgery makes the assessment of outcomes challenging. Key standards are described which should be adopted as a matter of priority. High risk procedures should be grouped and examined via HES data and a national audit of emergency laparotomy should be supported. The estimation of risk, timeliness of care at key points, the seniority of staff involved and the utilisation of critical care should be routinely recorded and reviewed.

Background

1. Introduction

The adult higher-risk non-cardiac surgical population represents a major healthcare challenge to every acute hospital. Surgery remains a common and effective treatment option for a diverse range of diseases and far from being replaced by drug therapies, surgery is now more frequently deemed a viable option for elderly patients and those with co-morbidities or advanced disease. The standard of patient care during surgery itself can now be extremely high and even complex elective surgery can be made relatively safe^{1 2}. However, successful surgery also depends on good peri-operative care and here lie challenges. Whilst we may have made some progress towards improving surgical outcomes, the available evidence suggests that post-operative adverse events may be much more frequent than many appreciate and that the consequences of these complications are considerable.

In the UK, the focus has fallen previously on cardiac surgery where specialist units carry out a modest range of predominantly elective procedures with routine intensive care support. Audit now shows good results which continue to improve with 2-3% mortality typical³. The established and transparent measurement of outcomes in cardiac surgery facilitates improvement by identifying centres of good practice and centres where change may be required.

By contrast, major general surgery is carried out in every acute hospital, encompassing a wide range of conditions which are, hence, more difficult to audit and conducted with limited critical care support. The mortality of elective major gastro-intestinal or vascular surgery substantially exceeds that of cardiac surgery. A much higher proportion of non-cardiac surgical patients are treated on an emergency basis and at present the service for such patients lacks focus despite much higher mortality and complication rates.

There is growing concern that this group of higher risk general surgical patients receive sub-optimal care which has important implications for patients and the healthcare economy. In the UK, 170,000 patients undergo higher-risk non-cardiac surgery each year⁴. Of these patients, 100,000 will develop significant complications resulting in over 25,000 deaths. General surgical emergency admissions are the largest group of all surgical admissions to United Kingdom hospitals and account for a large percentage of all surgical deaths. Emergency cases alone would account for 14,000 admissions to intensive care in England and Wales annually⁵. The mortality of these cases is over 25% and the ICU cost alone is at least £88 million.

Complications occur in as many as 50% of patients undergoing some common procedures, and these result in dramatic increases in length of stay and cost. Many of the patients undergoing this type of surgery are elderly with multiple co-morbidities^{6 7 8 9 10} and indeed the over 80s are more likely to present for emergency surgery than elective^{11 12}, where the risks multiply. Despite these findings there is surprisingly little research into how to improve these patients outcomes but structures of care which facilitate attention to the detail of peri-operative care, may help¹³.

Studies from the UK suggest that a readily identified higher risk sub-group accounts for over 80% of post-operative deaths but less than 15% of in-patient procedures^{4 6}. Advanced age, co-morbid disease, major and urgent surgery are the key factors associated with increased risk. Within this group, emergency major gastrointestinal surgery has one of the highest mortalities which can reach 50% in the over 80's⁸. Presently, this type of surgery is carried out in every acute hospital, but not always with consultant staff present and not always with routine admission to a critical care bed after surgery. Many of these issues were highlighted in the most recent NCEPOD report¹⁴

In the UK, fewer than one third of high-risk non-cardiac surgical patients may be admitted to critical care following surgery^{4 6}. In addition, those patients who do receive this level of care are discharged after a median stay of only 24 hours despite going on to have prolonged hospital stays. Premature discharge

from critical care has been identified as an important risk factor for post-operative death, as has delayed admission to critical care¹⁵. International comparisons suggest that critical care beds may run at 50% of comparable levels elsewhere, and indeed rank amongst the lowest in the developed world¹⁶.

To identify and advise on how these patients could be better managed, a joint working group was set up between the Royal College of Surgeons of England and the Department of Health to address these issues as they relate to the peri-operative care of general and vascular surgery in the first instance.

The following document seeks to explain to Commissioners, Chief Executives and Medical Directors the nature of the problem and to lay out logical steps which should be taken in order to achieve the greatest benefit in the most effective way.

2. Variation in current outcomes

There are several indicators that the outcomes from higher risk surgery in the UK are not as good as they should be. Review of 2008/9 HES data from Dr. Foster reveal a greater than 2-fold variation in relative risk of 30 day mortality (risk-adjusted) after non-elective lower gastrointestinal procedures between Trusts in The North West SHA. It is known that the chance of a patient dying in a UK hospital is 10% higher if they are admitted at a weekend rather than during the week¹⁷. There are no evident reasons for these differences other than that care, at times, is of variable quality: a conclusion which fits with the available evidence and professional opinion. International studies have reached similar conclusions and local audit data confirm that outcomes deteriorate if patients are admitted towards the end of duty periods and at weekends. Two recent NCEPOD reports, showed significant deficiencies in the active care of patients who ultimately died^{14, 18}. These included delays in assessment, decision making and treatment. There were shortfalls in access to theatre, radiology and critical care; surgery was suboptimally supervised in

30% of cases and there was a failure for juniors to call for help in 21% of cases. Timely surgery was not carried out in 22% of patients who died. There was also the failure to reliably administer therapy known to be of benefit such as antibiotic and venous thrombo-embolism prophylaxis. There are few data which compare our outcomes in the UK to other countries but one study reported that risk adjusted mortality rates were as much as 4 times higher in the UK than in the US¹⁹. A large percentage of the patients that survive have prolonged hospital stays with significant cost implications, both physical and emotional to the patient and their family, and financial to the hospital²⁰.

Together these data show that these higher-risk patients are a significant clinical burden in every hospital, use substantial Critical Care facilities with corresponding high cost but with outcomes which vary considerably between sites and within sites at weekends. These observations represent a poorly defined care pathway with standards that are either not determined or not implemented. The consequent impact on both patient outcomes and use of NHS resources is considerable. The scope for improvement is difficult to document given the very limited nature of current audit methods and the diversity of procedures undertaken. However, the findings are well recognised by many working in the field and nor are they surprising. Provision of services, particularly of theatre access, critical care and interventional radiology, is often incomplete and the correct location of patients after surgery is often not given sufficient priority. Furthermore, the clinical response for patients who deteriorate is often poorly thought through and at times, ad hoc. Aligning patients needs and subsequent risk of deterioration to the most appropriate pre and post-operative clinical area requires active early assessment of risk of death and clear objectives for clinical care to be identified.

3. How do adverse outcomes occur for the higher-risk general surgery patient?

While occasional patients die from haemorrhagic or cardiac complications during surgery, post-operative complications account for the bulk of morbidity

and mortality in general surgery. Some of these result from suboptimal surgical peri-operative care – perhaps on account of poor pre-operative preparation or inexperienced or delayed surgery or anaesthesia. For others, post-operative complications are chance occurrences but nevertheless ones which can often be readily anticipated and mitigated through consideration of co-existent diseases and the surgery performed. In the elderly, frailty is a risk factor and should be formally assessed in addition to nutritional and mental state¹⁴, Complications can be greatly reduced by optimal peri-operative care.

There are opportunities to improve outcomes before, during and after surgery. Many of these higher risk patients are emergencies where the time for pre-operative assessment is less and surgery is often unavoidable. In these cases, optimal resuscitation is important but delay is detrimental. However, for those patients undergoing elective high-risk surgery, optimal multidisciplinary pre-operative planning is the ideal.

Complications are common and raise costs, often several-fold. Their development reduces survival (short and long term) independently of pre-op risk and complexity of surgery²¹. Those that occur are managed variably and adverse outcomes are estimated to be due to errors in the process of care or medical management, each in about 20% of cases²².

Minor complications are extremely common after complex procedures and slow or suboptimal management of these, particularly in patients with other medical diseases can trigger a subsequent cascade of more serious complications. Many of the life threatening problems involve systemic infection (sepsis). Once a patient develops major complications, they are at risk of major organ dysfunction or failure. Typically, patients at risk or with organ dysfunction are managed in high dependency units (level 2), where the mortality is at least 5%. Once organ failure develops, full intensive care (level 3) is required and the mortality rises to 30% or more, often after prolonged treatment. The health and financial advantages of managing complex patients with adequate critical care support from the time of surgery are self evident.

Complications may be inevitable after this magnitude of surgery but their number and severity can be mitigated by rapid and successful treatment. It is well established that this requires the following steps

1. Rapid identification
2. Adequate resuscitation
3. Investigation to define the underlying problem
4. Rapid definitive treatment of that problem
5. Appropriate critical care provision to prevent further complications

Too often the whole process is slow or inaccurate as it is complex, requires multidisciplinary input, often occurs out of hours and is initiated by junior staff. Suboptimal care on general wards prior to critical care admission has been recognised as a cause of avoidable mortality¹⁵ which has resulted in the publication of a clinical guideline document from the National Institute for Clinical Excellence²³ and of a competency framework from DH²⁴. These documents outline a graded response strategy that each acute hospital should establish to recognise and respond to the deteriorating patient. Escalation of care for those that require surgical intervention, including radiological intervention, has not been the subject of specific guidance to date. Certainly in the US, the ability of different hospitals to manage complications differed significantly and this (rather than the initial frequency of complications) accounted for large variations in outcomes²⁵. Prompt intervention is fundamental to the successful treatment of the patient who deteriorates after surgery.

4. Sepsis

Sepsis (the body's generalised response to infection) requires special consideration because it is the principal reason for prolonged admission to critical care and death in these patients and because the existing guidelines

do not take into account current understanding of the timeliness of intervention.

The process is time critical and two steps are of particular importance. The first is to administer antibiotics within six hours as defined in the Surviving Sepsis Campaign²⁶.

The second is to deal with the source of sepsis which, in surgical practice, often means a complex operation or radiological drainage. Previous guidance with regard to the urgency of emergency surgery is too non-specific and does not take account of new evidence which suggests that patients with septic shock requiring source control have a progressive deterioration in outcome associated with increasing delay to source control²⁷. Delay of more than twelve hours after the onset of septic shock may increase mortality by a factor of 2.5 times when compared with patients who received source control within three hours. Gathering data on these patients is difficult but this expert group believes there is enough evidence at present to establish pragmatic guidance consistent with NICE CG50. Namely, that a graded response be established that requires increasingly rapid intervention for patients with increasing severity of illness and that the degree of urgency should be considerably greater than that previously accepted.

It is anticipated that the effects of this will be to reduce severity of illness, the need for higher levels of critical care and its associated cost and improve outcomes.

Actions

1. Managing the critically ill surgical patient with sepsis

Surgical patients may become critically ill for two reasons. They may present as an emergency with an acute surgical illness or they may develop complications following surgery or during surgical illness. Some complications have well defined treatment protocols and others are so catastrophic that the need for immediate summoning of the cardiac arrest team is obvious. However, the graded response for identification and treatment of sepsis, the most frequent serious complication is not well defined. This deficit leads to avoidable adverse outcomes.

(i) Escalation of care

Fundamental to prompt definitive treatment of sepsis and indeed, all complications, is the need to identify critically ill patients at an early stage. This escalation guideline is written with reference to existing documents; NICE CG50²³ and Competencies for Recognising and Responding to Acutely Ill Patients in Hospital²⁴. The graded response to early warning scores will be described as a three point scale of response to low, medium and high scoring patients.

Surgical patients frequently differ from non-surgical ones in two ways. Firstly, the conditions which develop often demand greater urgency and secondly, they more often require complex operative interventions following advanced imaging. These differences bring opportunity for delay.

For a medium-score patient NICE CG50 requires: "Urgent call to team with primary medical responsibility and simultaneous call to staff with core competencies in care of acute illness". In the case of a surgical patient that has deteriorated on the ward the member of staff with "core competencies" is

a surgical trainee, who will usually have passed MRCS. A typical “medium score” patient would be one that is developing severe sepsis or one with less severe acute pathology but with significant co-morbidities.

This trainee, here denoted MRCS, is the secondary responder in the chain of response described ²³. The MRCS plays a key role in diagnosis and communication between tertiary response groups; crucially the consultant surgeon although microbiologist, radiologist, anaesthetist and intensivist may all need to be involved within a short space of time. Staffing arrangements between hospitals will vary. Responsibility for ensuring that the MRCS is able to review a patient that triggers a medium score without delay is fundamental and will rest with individual departments.

For the escalation structure, below, to work for the patient’s benefit, the MRCS must be competent in recognising whether a deteriorating patient has sepsis or not and whether the cause of sepsis is most appropriately treated with antibiotics alone or with source control. The MRCS must also be able to differentiate between the different levels of severity of sepsis. Successful attendance at a CCrISP Course ²⁸, or equivalent would provide this and this is a “strongly recommended” facet of basic surgical training in the UK.

Suggested pathways for escalation are shown in the appendices. That in appendix 2 utilises the early stages of the generic pathway described in NICE CG50 up to the point of referral to the secondary responder. However note that 12 hourly observations is too infrequent for this group: hourly observations would be more usual until medical review and would likely be triggered by the Early Warning Score (EWS). There follows the recommended pathway for the surgical patient. Further explanation of the current status of Early Warning Scores is given in Appendix 1.

The summary timelines for assessment of the unstable patient and for intervention are shown below. For definitive treatment to occur within the recommended timeframe, it will be clear that each phase of treatment must be

expeditious. These phases often include initial recognition, initial assessment, MRCS assessment, investigation (most commonly CT scan) and senior decision making. Hospitals should audit the stages of the pathway to minimise the avoidable delays which are currently recognised.

(ii) Urgency of source control

Patients with sepsis require immediate broad-spectrum antibiotics with fluid resuscitation and source control

- a) Those with septic shock require immediate broad-spectrum antibiotics with fluid resuscitation and source control. Delay to source control of more than twelve hours after onset of hypotension when compared with a delay of less than three hours could be expected to increase mortality from 25% to more than 60%²⁷. Rapid involvement of senior staff is important. Control of the source of sepsis by surgery or other means should be immediate and underway within three hours.
- b) Patients with severe sepsis (sepsis with organ dysfunction) are at greatest risk of developing septic shock. There is no direct evidence to confirm that delayed source control worsens outcome but there are obvious advantages to operating before progression to septic shock occurs^{29 30 31} given the associated 5 to 10 fold rise in mortality which occurs as the patient deteriorates. Surgery or equivalent (e.g. radiological drainage) should be carried out within 6 hours from the onset of deterioration. These patients require immediate broad-spectrum antibiotics with fluid resuscitation, urgent but not immediate surgery, frequent monitoring (as per CG50) in an appropriate environment during the interim to promptly identify development of hypotension. Where it is elected to observe and resuscitate the patient for a few hours until morning, surgery should assume priority over elective procedures. Neither observation nor resuscitation should delay source control for more than 6 hours. Evidence suggests that further delays at this point are common^{14 32}.

- c) Source control for patients with sepsis but without organ dysfunction should always be carried out within 18 hours. Immediate broad-spectrum antibiotics are required but surgery can be delayed overnight or until the next theatre becomes available. Source control is needed before progression to severe sepsis which carries a greater overall mortality and an increased frequency of observations is needed in the interim to identify any clinical deterioration which should trigger a revised management plan.
- d) Patients that require source control but have not mounted a systemic inflammatory response are clinically appropriate for NCEPOD classification "Expedited".

Doctors should be aware of these timescales when determining the urgency of assessment and intervention. As the acute management pathway for many of these patients is tortuous (assessment, senior assessment, investigation, anaesthetic review, critical care review, theatre scheduling, operation), the need for urgency at each stage is emphasised.

These timescales shown are the maximum. Some patients will have surgical considerations mandating more urgent intervention.

Hospitals should provide adequate emergency theatre access free from predictable obstruction or restriction caused by over-running elective work or manpower shortage. This is not infrequently seen at late afternoon / early evening.

Hospitals should also ensure that there are clear arrangements in place for interventional radiology, especially out of hours. For many, this will be via a network of cover across multiple hospitals.

Moving a patient to critical care does not treat the source of sepsis and the focus must remain on timely definitive care. This needs to be balanced with appropriate but rapid pre-operative resuscitation. If the patient becomes

hypotensive, fails to respond to resuscitation or otherwise deteriorates then immediate treatment is necessary as in a).

(iii) Summary timelines

a) Surgical Response (level 2 / secondary)

| <i>EWS</i> | <i>Grade of staff</i> | <i>time</i> |
|-------------------|--|--------------------|
| Low | Foundation / ST 1-2 | 1 hour |
| Medium | MRCS | within 30 mins |
| High | MRCS and critical care / anaesthetic staff | immediate |

If there is an incomplete response to resuscitation within 1 hour, particularly if remains hypotensive or with organ dysfunction then : Inform / involve senior and move to critical care area or operating room as appropriate.

If MRCS not available because operating, the ICU or anaesthetic SpR should be called directly to the patient and the consultant surgeon involved. At each stage, all members of the multidisciplinary team should be encouraged to involve more senior staff if there is a delayed or incomplete response by the medical team or the patient.

b) Intervention to control source of sepsis

| <i>Severity of sepsis</i> | <i>Time to intervention (maximum)</i> |
|-----------------------------------|--|
| Septic shock | Immediate |
| Severe sepsis / organ dysfunction | as soon as possible and within 6 hours of onset |
| Sepsis | as soon as possible and within 18 hours (7am – 10pm start) |
| Infected source, no SIRS | as soon as possible (7am – 10pm start) |

2. Assessing and identifying risk

Why it should be done

Studies from the UK suggest that a readily identified higher-risk sub-group accounts for over 80% of post-operative deaths but less than 15% of in-patient procedures^{4 6}. Advanced age, co-morbid disease, major and urgent surgery, primary diagnosis and acute physiological deterioration are the key factors associated with increased risk. Routine identification of patients most at risk would permit care and resources to be better directed.

How should risk be assessed?

Presently, clinicians' assessment of peri-operative risk may be omitted, inaccurate or may not lead to an effective change in clinical management. Objective assessment of risk must become routine. Most importantly, identification of higher risk needs to trigger joint advance planning specific to that case.

1. We recommend that objective risk assessment become a mandatory part of the pre-operative checklist to be discussed between surgeon and anaesthetist for all patients. This must be more detailed than simply noting the ASA score.
2. For elective patients, risk should be assessed at pre-operative assessment and those at high risk should see the anaesthetist who will anaesthetise them. On occasion being seen by a consultant colleague from a small specialist team working to the same agreed protocol would be acceptable. A range of risk scores and tests of exercise capacity are available and should be adopted. Close working arrangements and subspecialisation is advocated for higher risk cases.

Patients with a predicted mortality $\geq 10\%$ should be managed as "high risk". Approximately 20% of general surgical emergency procedures fall in this

category together with complex elective GI and vascular procedures, in comorbid patients.

There are a number of methods with which to predict hospital mortality risk. Some methods are described below. Each method has strengths and weaknesses so for patients to be safely defined as low risk they should not obviously exceed high risk criteria for any method.

Note that the average mortality of a defined group can be expected to be approximately 2 to 4 times the threshold and it is anticipated that teams may wish to set the threshold lower in time (5%). An estimated risk of 5 to 10% may usefully define a medium risk group.

- a) P-POSSUM, freely available on the internet³³ is possibly the simplest and best validated method and a good place to start. Its scoring includes operative details so these have to be estimated for pre-operative use and can be updated at the end of surgery.

- b) Alternatively, the criteria below are taken from an expert clinical trial in this population and also fit with expert opinion. These will define a group with a predicted mortality >5% and an overall mortality of 10-12%.

Patients undergoing major gastro-intestinal or vascular surgery who are either:

1. aged >50 years
 - a. and undergoing urgent, emergency or re-do surgery
 - b. or have acute or chronic renal impairment (serum creatinine >130 $\mu\text{mol/l}$)
 - c. or have diabetes mellitus (even if only diet controlled)
 - d. or have or are strongly suspected clinically to have any significant risk factor for cardiac or respiratory disease

2. are aged >65 years

3. have shock of any cause, any age group.

- c) A third way of identifying the higher risk surgical patient is by reference to Hospital Episode Statistics (HES) procedure groups. While this approach shows considerable concordance with the methods above for populations of patients, its failure to include acute illness or chronic co-morbid disease means it should be used alongside a consideration of patient physiology for individual patient assessment. With that caveat, HES data analysis shows that the following emergency cases have an average mortality of >10% in the UK; laparotomy for peritonitis, resection of colon or rectum, therapeutic operations on small bowel, therapeutic upper GI endoscopy, peptic ulcer surgery, gastrectomy and splenectomy. In such cases patients are likely to be "higher risk" unless they are unusually fit.

Other physiological derangements, disease states and procedures may also define high and medium risk patients, including urgent surgery in patient with ASA ≥ 3 + at least 1 acute organ dysfunction/failure, ASA 4 or 5, dialysis dependent patients or patients with elevated lactate (>4 mmol/l).

The identification of higher risk status should lead to certain levels of care. Staff involved should be sufficient in seniority and number to permit care to proceed expeditiously. It is recognised that while some more senior trainees may have many of the skills necessary, this is less so than previously. Furthermore, the presence of a consultant can remove organisational barriers and assist in prompt decision making. For the surgical team, this practical assistance is essential given modern day on call arrangements. Anaesthetic juniors may similarly lack experience and have to manage calls about other patients simultaneously, causing further delays. Consequently, the starting position should be that each higher risk case (predicted mortality $\geq 10\%$) should have the active input of consultant surgeon and consultant anaesthetist in theatre. Occasional cases may be appropriately managed by unsupervised juniors but this should be an active and audited senior decision. Calling senior staff at a later stage once problems have developed will usually be associated with worse outcomes and this event should also be audited. It is also recognised that the systemic impact of sepsis on patients undergoing

major procedures is not always identified initially and seniors should be cautious about leaving before the case is finished.

End of Surgery Bundle

The post-operative pathway must be determined by the risk of death and complications and receiving areas must possess the competencies to deal with surgical patients.

A key decision point occurs towards the end of higher-risk surgery, much of which is emergency in nature and thus less than perfectly planned. At this point, decisions need to be made concerning the disposition of the patient following surgery. Underestimating the degree of existing physiological upset or of the likely evolution of organ dysfunction can be catastrophic: late admission to critical care carries a much higher mortality than a planned admission from the operating room. Staff may be junior, tired or dealing with a relatively unfamiliar set of circumstances and it seems logical to conduct a structured assessment of risk towards the end of surgery. One method would be to use the Apgar score for surgery³⁴. An alternative would be to use the bundle described below³⁵ within the last 30 minutes of surgery in all cases identified by the pre-operative assessment of mortality risk $\geq 5\%$ and in those who deteriorate during surgery.

1. Risk score patient ($\geq 10\%$ mortality defines high risk)
2. Check Arterial Blood Gases to assess lactate, acid-base status and the ratio of arterial oxygen concentration to the fraction of inspired oxygen (P:F ratio)
3. Summarise fluids given and draft ongoing fluid requirements.
4. Reverse muscle relaxant; use of nerve stimulator is mandatory.
5. Check and document temperature, plan further correction as necessary.

Based on the bundle criteria, the surgeon and anaesthetist should decide jointly the preferred destination of the patient after surgery. All high risk patients should be admitted to the appropriate (level 2 / 3) critical care unit

with surgical competencies. This decision will be influenced by adverse events during surgery or a high likelihood of deterioration in the short to medium term. The bundle should be used to supplement rather than replace existing indicators of the need for critical care.

1. The POSSUM score is the most validated risk prediction method for general and vascular patients that takes into account pre-operative and peri-operative factors. P-POSSUM may be used for all patients³⁶. A predicted mortality risk $\geq 10\%$ indicates need for critical care admission, except for patients on End of Life pathways with appropriate palliative care facilities available at ward level.
2. Hyperlactataemia (>4 mmol/l) and significant metabolic acidosis indicate unresolved physiological impairment that requires ongoing invasive monitoring +/- physiological support³⁷. Serum lactate levels may also be used to guide fluid therapy and levels >2 mmol/l indicate the need for closer monitoring³⁸. P:F ratio < 40 kPa is consistent with an acute lung injury. A senior critical care specialist should be involved in the decision to extubate. A P:F ratio <26 kPa is consistent with a diagnosis of ARDS: the patient should be transferred to ICU intubated.
3. Both excessive and inadequate intravenous fluid administered in the peri-operative and postoperative period can be harmful particularly in the elderly¹⁴. A fluid plan should be agreed between the anaesthetic team and senior surgeon, bearing in mind modern guidelines and the risks of both excessive and inadequate fluid therapy³⁹. This should include blood loss and replacement.
4. Partial reversal of muscle relaxation is common and poorly recognised. It is a risk factor for post-operative respiratory failure and aspiration. Nerve stimulation and reversal is mandatory if a neuromuscular blocker has been given regardless of time interval. A Train-of-four (TOF) ratio of 0.9 is required for airway protection. Unfortunately TOF ratio is difficult to assess

accurately by observation alone⁴⁰. To be confident of airway protection, neostigmine should not be given if the TOF count is less than 2 and at least 9 minutes should elapse after neostigmine bolus before extubation is attempted.

5. Hypothermia (core temperature $<36^{\circ}\text{C}$) increases the incidence of post-operative myocardial events⁴¹ and wound infections. Drug metabolism is reduced such that duration of neuromuscular blockers can be doubled⁴² and neostigmine can take 20% longer to take effect⁴³. NICE clinical guidance (Management of Inadvertent Perioperative Hypothermia, 2008) should be followed⁴⁴.

The use of “bundles” has been shown to increase the reliability of key steps of care⁴⁵. The concept of using a bundle at the end of high risk surgery should be tested in individual institutions, if necessary adjusted for context, and if found to increase the reliability of key step delivery, incorporated into routine anaesthetic paperwork. Joint early discussion with the critical care team is fundamental.

Postoperative care

Access to critical care is an essential aspect of adequate peri-operative care for the high-risk group in order to identify complications early and minimise their impact.

All patients should be managed after surgery in a location determined by risk and staff competence. Hospitals should plan their critical care resource to match need in order to avoid shortages and define critical care areas accordingly.

Patients should move up and down through a spectrum of levels of care. Levels of care are described as⁴⁶:

| Level | Description | Patient characteristics |
|-------|-----------------|--|
| 0 | ward | |
| 1 | enhanced ward | At risk of deterioration |
| 2 | high dependency | Needs detailed observation, intervention or single organ support |
| 3 | intensive care | Multiple organ support, complexity |

All patients with a predicted mortality of $\geq 10\%$ should be admitted to a level 2 or 3 critical care area after surgery and all patients should have an updated management plan which incorporates haemodynamic and blood gas parameters, on-going antibiotics, nutrition and thromboembolic prophylaxis.

Trusts may wish to examine their existing provision particularly around levels 1 and 2. When compared to Level 0 care, the impact of Level 1 or 2 care is likely to be much greater in the unscheduled surgical population than the elective population due to the dynamic nature of the acute illness and its influence on organ function. Recognition of any deterioration in organ function and timely intervention is essential to optimise patient benefit. Provision of this level of monitoring is frequently difficult to deliver in a standard ward environment with staffing ratios per patient which is frequently < 0.20 nurse : patient. Defining pathways for such patients affords organisations an opportunity to address competencies of staff and staffing ratios to deliver a tiered pathway of care.

Some organisations have developed bespoke solutions to this such as the development of PACUs or co-locating medium risk patients in pre-defined clinical areas.

1. Structured care on the Post anaesthetic care unit (PACU)

A patient inappropriate for the ward could be admitted to PACU for continued monitoring. Formal joint assessment should occur after four hours. If the patient is alert and has a normal temperature, mean arterial pressure, pH, lactate and gas exchange, and the previous three consecutive hourly urine

volumes were all > 0.5 ml/kg, transfer to the ward is acceptable unless there is specific clinical concern to the contrary.

If the above criteria are not met after four hours in PACU, care should be formally taken over by the critical care team who will continue to care for the patient in PACU until transfer to a critical care bed can be arranged or the patient is considered ready for transfer to the ward by a senior critical care specialist.

To do this, hospitals will need to ensure that there is a 24/7 PACU service and that a consultant from anaesthesia/ critical care / surgery is identified to take responsibility for this provision and to work with the PACU manager to ensure delivery of appropriate care.

Ongoing audit will allow assessment of impact on PACU and elective surgery. Hospitals will wish to make the difference between PACU and theatre recovery explicit as inadequate staffing may result in loss of ability to undertake further emergency surgery if a patient is “blocking” recovery. These events should be audited and classified as an adverse incident.

2. Co-location of medium risk patients

Existing systems of critical care can leave a large step between HDU and ward care. The co-location of medium risk patients in special wards or ward-areas (level 1) could be expected to lead to immediate improvement in standards even if staffed near general surgical ward levels and without significant investment in additional monitoring.

Immediate benefits would be promoted provided Trusts:

- Establish local protocols drawn up jointly between surgical and critical care departments to define parameters of care and to ensure seamless transition of patients between units
- Establish co-operative education programmes with critical care for nursing and medical staff
- Establish improved daily communication between units

- Recommend geographical proximity to critical care where possible
- Name a critical care consultant with responsibility for basic education and support for nursing and junior medical staff.

Audit and Outcomes

The relative paucity of data in this field needs to be addressed urgently, preferably on a national basis. Given the mortality and morbidity associated with this group, comparative risk-adjusted outcomes should be monitored for each hospital. At the moment, HES data may be the best available. The adoption of a defined basket of HRG codes would facilitate this. International comparisons would provide the greatest re-assurance that care for this group is optimal.

The processes advocated in this report should be audited in each hospital and key indicators include:

- Outcomes (death, length of stay) from higher risk general surgery
- Frequency of observations in higher risk group
- Accuracy of risk estimate prior to surgery
- Accuracy of risk estimate at end of surgery
- Time to CT from emergency admission or deterioration
- Time from deterioration to operation for higher risk group
- Compliance with the standard for intra-operative surgical team seniority
- Compliance with post-surgery pathway for higher risk patients.
- Unplanned Surgical readmissions to Critical Care within 48hrs of discharge back to the ward.

Emergency laparotomy is a clearly defined point in the pathway of a significant proportion of these patients and in this group, many of the factors discussed in this report come together. The Laparotomy network audit <http://www.networks.nhs.uk/nhs-networks/emergency-laparotomy-network> is beginning to look at these patients on a voluntary basis and this study should be supported and expanded.

Conclusions

Peri-operative care of higher risk general surgical patients in the UK appears to have significant deficiencies. Outcomes are variable, appear worse than other countries and generate a large health cost through prolonged hospital stay and use of intensive care.

While there are several specific initiatives (e.g. Hospital Acquired Thrombosis) and patient pathways for single operations (e.g. Aortic Aneurysm), there is a lack of overall recognition and strategy for the care of patients at higher risk of death and complications.

Standards of care are described in this document. Trusts should develop pathways in order to achieve these.

This higher risk group comprises 12 to 15% of cases but contributes 80% or more of postoperative deaths and complications. This group can be identified at an early point and differential management pathways applied. Identification of these at risk patients should become a formal part of patient assessment and included in the pre-operative checklist.

An estimated mortality of $\geq 10\%$ defines a high risk patient. An estimated mortality of $\geq 5\%$ defines medium risk. Together they can be termed “higher-risk”.

In particular, attention could be better focussed on elective cases who develop complications and on major emergency cases. A defined and escalating pathway of management, which complements existing guidance for acute care, should be adopted. The described pathways match urgency to patient need and include guidance on senior involvement and time to treatment.

The principal life threatening complication is the development of severe sepsis. Patients from this group account for the greatest use of ICU beds.

Improved assessment and treatment would likely improve outcomes and reduce ICU utilisation.

High risk procedures should be managed by consultant staff.

There should be a brief but structured review of progress towards the end of higher risk operations, conducted jointly between surgeon and anaesthetist. This End of Surgery bundle should guide the location of post-operative care.

High risk patients should be managed after surgery in a level 2 or 3 critical care area. There appears to be a shortfall of critical care beds which Trusts should address. The format of these needs consideration in order to find the most effective and cost efficient structures as several different models of care exist. We believe that investment in better perioperative care would realise benefits for both cost and outcomes.

Outcomes from emergency surgery are difficult to compare due to the range of diagnoses and operations. A national audit of higher-risk emergency surgery is essential. A basket of HES codes is proposed and should be agreed for ongoing comparison.

Appendix 1 Early Warning Score (EWS).

This is a scoring system used in physiological track and trigger systems. The score is based on routinely recorded physiological observations such as blood pressure, heart rate etc. Each observation is given a score of zero if it is normal increasing to (typically) 3 as the observation deviates further from the normal range. The sum of all parameter scores gives a total EWS. There is currently no national system in use. Different hospitals use scoring systems that differ in the methodology for generating the final EWS and in the response system. Until a national system is established (and audited) the only generally applicable guidance comes from NICE CG50 which stipulates that hospitals should establish a graded response system according to the following system:

Low-score group: increase frequency of observations and inform nurse in charge.

Medium-score group: urgent call to team with primary medical responsibility and simultaneous call to personnel with core competencies for acute illness.

High-score group: emergency call to team with critical care competencies and diagnostic skills.

It is for individual trusts to determine what EWS score triggers each of these responses.

Septic Shock is defined as severe sepsis complicated by persistent hypotension (systolic less than 90mmHg or >40% decrease from baseline) that is not reversed by fluid resuscitation. An adequate volume of fluid is considered to be 20ml/kg of crystalloid or an equivalent volume of colloid. In this document hypotension in the context of severe sepsis is taken to be persistent hypotension that is not fluid responsive.

Appendix 2.


Pathway for assessment and response for unwell surgical patient

Figure 1, below, combines initial generic assessment taken from Nice CG50 (upper part of figure) with a surgery specific pathway (lower part of figure). Initial routine monitoring for this group of patients will be hourly.

Figure 1

1.2.3 Care pathway

Assessment and monitoring



MRCS to attend patient and to coordinate response.

MRCS will immediately leave less urgent tasks such as clinics and ward rounds and will delegate to an appropriately competent colleague if currently operating or attending another medium-high score case.

Is the cause of deterioration medical or surgical?
Needs MRCS+ input to this decision

Medical

Surgical

Continue to follow NICE CG50

Immediate life, limb or organ saving surgery is indicated

Resuscitation is simultaneous with intervention. Example; the exsanguinating patient. MRCS to liaise with consultant surgeon, anaesthetist and theatre staff. The patient should be transferred to theatre within minutes of the decision to operate.

The patient is septic

The need for source control must be established rapidly. Urgency of surgery depends on severity of sepsis.

The patient has sepsis but no organ impairment or low score risk. *Establish source control urgently and always within 18 hours. Patient should be monitored hourly and reassessed by MRCS every 6 hours to check for progression to severe sepsis/septic shock.*

The patient has severe sepsis or medium-high score risk without hypotension. *Establish source control as soon as possible and within 6 hours maximum. Reassess hourly for progression to septic shock and provide appropriate interim critical care.*

The patient has septic shock. *The patient's chance of survival progressively deteriorates with increasing delay to source control. Establish source control as soon as possible. Transfer to theatre must not be delayed for resuscitation which should be continued in the anaesthetic room.*

The patient is NOT septic and does not require immediate intervention

Organise initial treatment and investigations, liaise with consultant surgeon and plan definitive treatment.

Appendix 3

Draft pathway for unscheduled admissions

References

-
- ¹ Jones RS: Comparative mortality in anaesthesia. *Br J Anaesth* 2001, 87(6):813-815.
- ² Haynes AB, Weiser TG, Berry WR, et al: A surgical safety checklist to reduce morbidity and mortality in a global population. *N Engl J Med* 2009, 360(5):491-499.
- ³ <http://heartsurgery.cqc.org.uk/Survival.aspx>
- ⁴ Jhanji S, Thomas B, Ely A, Watson D, Hinds CJ, Pearse RM: Mortality and utilisation of critical care resources amongst high-risk surgical patients in a large NHS trust. *Anaesthesia* 2008, 63(7):695-700.
- ⁵ Intensive Care National Audit & Research Centre (ICNARC), London 2010. Data derived from Case Mix Programme Database based on 170,105 admissions to 185 adult, general critical care units in NHS hospitals across England, Wales and Northern Ireland.
- ⁶ Pearse RM, Harrison DA, James P et al. Identification and characterisation of the high-risk surgical population in the United Kingdom. *Crit Care* 2006, 10(3):R81.
- ⁷ Cullinane M, Gray AJ, Hargraves CM, et al, : The 2003 Report of the National Confidential Enquiry into Peri-Operative Deaths. In. London: NCEPOD; 2003.
- ⁸ Association of Surgeons of Great Britain and Ireland. Emergency General Surgery: The future. A consensus statement. June 2007. Available from: http://asgbi.org.uk/en/publications/consensus_statements.cfm
- ⁹ Semmens JB, Aitken RJ, Sanfilippo FM, et AL. The Western Australian Audit of Surgical Mortality: advancing surgical accountability. *Med J Australia* 2005; 183 (10):504-508.
- ¹⁰ Cook TM, Day CJE. Hospital mortality after urgent and emergency laparotomy in patients aged 65yr and over. Risk and prediction of risk using multiple regression analysis. *Brit J Anaes* 1998; 80; 776-781.
- ¹¹ Ford PNR, Thomas I, Cook TM, Whitley E, Peden CJ. Outcome in critically ill octogenarians after surgery: an observational study. *British Journal of Anaesthesia* 2007; 99(6):824-829.
- ¹² Scottish Audit of Surgical Mortality Report 2008 (2007 data) www.sasm.org.uk
- ¹³ de Vries E, Prins H, Crolla R, Effect of a Comprehensive Surgical Safety System on Patient Outcomes. *N Engl J Med* 2010;363:1928-37.
- ¹⁴ Elective & Emergency Surgery in the Elderly: An Age Old Problem (2010) (http://www.ncepod.org.uk/2010eese.htm?utm_source=Sign-Up.to&utm_medium=email&utm_campaign=220074-NHS+Institute+Alert+-+December+2010)
- ¹⁵ McQuillan P, et al. Confidential inquiry into quality of care before admission to intensive care. *BMJ*.1998;316(7148):1853-9.
- ¹⁶ Adhikari NKJ, Fowler RA, Bhagwanjee S, et al. Critical care and the global burden of critical illness in adults. *Lancet* 2010;376:1339-46.
- ¹⁷ Aylin P, Yunus A, Bottle A, et al. Weekend mortality for emergency admissions. A large, multicentre study. *Qual Saf Health Care*. 2010; 19:213-7.
- ¹⁸ National Confidential Enquiry into Patient Outcome and Death. 2009 Deaths in Acute Hospitals: Caring to the End? London. http://www.ncepod.org.uk/2009report2/Downloads/DAH_report.pdf
- ¹⁹ Bennett-Guerrero E, Hyam JA, Shaefi S et al. Comparison of P-POSSUM risk-adjusted mortality rates after surgery between patients in the USA and the UK. *Br J Surg*. 2003;90:1593-8.

-
- ²⁰ Clarke PA, Murdoch H, Thomas MJ, Cook TM, Peden CJ. Mortality and postoperative care after emergency laparotomy. *Euro J Anaesthesiol*. 2010;[epub ahead of print].
- ²¹ Khuri SF, Henderson WG, De Palma RG et al. Determinants of long-term survival after major surgery and the adverse effect of postoperative complications. *Ann Surg*. 2005 Sep;242(3):326-41
- ²² Itani KM, Denwood R, Schiffner T et al. Causes of high mortality in colorectal surgery: a review of episodes of care in Veterans Affairs hospitals. *Am J Surg*. 2007 Nov;194(5):639-45.
- ²³ NICE clinical guideline 50 – Acutely ill patients in hospital. NICE, 2007.
- ²⁴ Competencies for Recognising and Responding to Acutely Ill Patients in Hospital. Department of Health, 2009
- ²⁵ Ghaferi AA, Birkmeyer JD, Dimick JB: Variation in hospital mortality associated with inpatient surgery. *N Engl J Med* 2009, **361**(14):1368-1375
- ²⁶ Levy MM, Dellinger RP, Townsend SR, et al. Surviving Sepsis Campaign: results of an international guideline-based performance improvement program targeting severe sepsis. *Crit Care Med*. 2010;38:367-74.
- ²⁷ Outcome of Septic Shock Correlates with Duration of Hypotension Prior to Source Control Implementation. ICAAC, 2004
- ²⁸ Care of the Critically Ill Surgical Patient Course. Available around the country, organised by The Royal College of Surgeons of England (www.rcseng.ac.uk)
- ²⁹ Sundararajan V, Korman T, Macisaac C, et al. The microbiology and outcome of sepsis in Victoria, Australia. *Epidemiol Infect*. 2006 Apr;134(2):307-14
- ³⁰ Angus DC, Linde-Zwirble WT, Lidicker J, et al. Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated costs of care. *Crit Care Med*. 2001;**29**:1303–1310.
- ³¹ Annane D, Aegerter P, Jars-Guincestre MC, Guidet B. Current epidemiology of septic shock: the CUB-Rea Network. *Am J Respir Crit Care Med*. 2003;**168**:165–172.
- ³² ASGBI survey on Emergency Surgery, 2010. Association of Surgeons of GB & I, London, Sep 2010. <http://asgbi.org.uk/en/publications/newsletters.cfm>
- ³³ <http://www.riskprediction.org.uk/index.php>
- ³⁴ Gawande AA, Kwaan MR, Regenbogen SE. An Apgar Score for Surgery. *J Am Coll Surg* 2007;204:201–208
- ³⁵ Peden CJ. Improving outcome in high risk surgical patients. Practicum for Masters in Public Health (Clinical Effectiveness). Harvard School of Public Health 2009
- ³⁶ Prytherch DR, Whiteley MS, Higgins B, et al. POSSUM and Portsmouth POSSUM for predicting mortality. *Br J Surg* 1998; **85**: 1217–1220
- ³⁷ Dellinger RP, Levy MM, Carlet, JM, et al: Surviving Sepsis Campaign: International guidelines for management of severe sepsis and septic shock: 2008. *Crit Care Med* 2008; 36:296 –327
- ³⁸ Wenkui Y et al. Restricted peri-operative fluid administration adjusted by serum lactate level improved outcome after major elective surgery for gastrointestinal malignancy. *Surgery* 2010;147:542-52
- ³⁹ Powell-Tuck J, Gosling P, Lobo DN, et al. (2008) British consensus guidelines on intravenous fluid therapy for adult surgical patients. GIFTASUP. Available from http://asgbi.org.uk/en/publications/surgical_resources_and_documents.cfm

-
- ⁴⁰ Eriksson LI, Sundman E, Olsson R, et al. Functional assessment of the pharynx at rest and during swallowing in partially paralyzed humans: simultaneous videomanometry and mechanomyography of awake human volunteers. *Anesthesiology* 1997; 87:1035-43
- ⁴¹ Frank SM et al. Perioperative maintenance of normothermia reduces the incidence of morbid cardiac events: A randomised clinical trial. *JAMA* 1997; 277:1127-34
- ⁴² Heier T et al. Mild intraoperative hypothermia increases duration of action and spontaneous recovery of vecuronium blockade during nitrous oxide-isoflurane anaesthesia in humans. *Anesthesiology* 1991; 74:815-9
- ⁴³ Heier T et al. The influence of mild hypothermia on the pharmacokinetics and time course of action of neostigmine in anesthetised volunteers. *Anesthesiology* 2002; 97:90-5
- ⁴⁴ NICE Clinical Guidance 065: Management of Inadvertent Perioperative Hypothermia, 2008.
- ⁴⁵ Resar R, Pronovost P, Haraden C, Simmonds T, Rainey T, Nolan T. Using a bundle approach to improve ventilator care processes and reduce ventilator-associated pneumonia. *Joint Commission on Quality and Patient Safety*. 2005;31;243-248
- ⁴⁶ http://www.datadictionary.nhs.uk/data_dictionary/attributes/c/cou/critical_care_level_de.asp?shownav=1